

ENVIRONMENTAL HAZARD 8

Do not apply directly to water, areas where surface water is present or to intertidal areas below mean high water mark. Do not contaminate with water when cleaning equipment or disposing of equipment wash water. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Keep containers tightly closed when not in use. Keep away from heat and open flame. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not reuse empty containers. Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by the state authorities, by burning. If burned, stay out of smoke.

PROHIBITION AND WARRANTY

It is a violation of FPA rules to use this product in a manner inconsistent with its label. The directions, recommendation and other technical information given are for guidance. We make no warranty of any kind, expressed or implied concerning any damage or effect that fails to take place. User assumes all risks and liabilities resulting from handling, use or application.

STIMULATE

YIELD ENHANCER

ACTIVE INGREDIENTS

CYTOKININ	= 00.009%
AUXINS (IBA)	= 00.005%
GIBBERELLINS	= 00.005%

FPA REG. NO. 1 - 5IF - 043

NET CONTENT: 1 Liter

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing mist. May cause irritation of nose, throat, and/or skin. Avoid contact with eyes, skin and clothing. If swallowed: induce vomiting by sticking finger down throat. Never give anything by mouth to an unconscious person. Get medical attention. If in eyes: wash with plenty of clear water for at least 16 minutes. Get medical attention.

STIMULATE

GENERAL INFORMATION

STIMULATE™ Yield Enhancer is a biostimulant containing natural plant growth regulators and chelated trace elements. STIMULATE™ enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizers, but it is also compatible with pesticides.

FOLIAR APPLICATIONS (per hectare basis)

Use 250 ml STIMULATE™ diluted in sufficient amount of water and apply 14 days after transplanting. Repeat spray during panicle initiation using the same amount of STIMULATE.

SEED TREATMENT

Use 2.5 ml of STIMULATE diluted in water per kilogram of seeds.

Exclusively Distributed in the Philippines by:



STOLLER Philippines, Inc.
Unit 1001, Manila Luxury Condominium
Pearl Drive, Ortigas Center, Pasig City
1600 Philippines, Tels. 625-5289, 624-2121

A Product of:



STOLLER International, Inc.
Houston, Texas