

# Harvest More

30 - 10 - 10

NET. WT. 1 kg.

# Harvest More

30 - 10 - 10

WATER SOLUBLE FOLIAR FERTILIZER  
WITH TRACE ELEMENTS

FPA Reg. NO. 1-1 IF - 1145

## Harvest More

30 - 10 - 10

WATER SOLUBLE FOLIAR FERTILIZER WITH TRACE ELEMENTS

HARVEST MORE fertilizer is a highly nutritious plant food which can be absorbed through the leaves as well as the roots of the plant. It provides practically all the nutrients the plant needs in order to thrive.

### GUARANTEED ANALYSIS

Fertilizer Grade:  
Nitrogen (N) 30.0%  
Phosphate (P<sub>2</sub>O<sub>5</sub>) 10.0%  
Potassium 10.0%

### Trace Elements:

Zinc 1000 ppm  
Copper 500 ppm  
Manganese 500 ppm  
Boron 200 ppm  
Molybdenum 50 ppm  
Cobalt 20 ppm



**STORAGE & DISPOSAL:**  
Store in a cool and dry place away from food, feedstuffs and water sources. Dispose empty containers in a sanitary landfill or by burning. If being burned, keep away from smoke.

**COMPATIBILITY:**  
Compatible with most commonly use soil fertilizers and pesticides. If using previously untried mixtures, it is suggested to mix a small amount first before preparing a full tank mix.

**PROHIBITION & WARRANTY:**  
It is a violation of FPA rules to use this product in a manner inconsistent with its label. The directions, recommendation and other technical information given for guidance. We make no warranty of any kind, expressed or implied concerning any damage or effect that fails to take place. User assumes all risks and liabilities resulting from handling, use or application.

Manufactured in the USA by:  
**STOLLER**  
International, Inc.  
Houston, Texas

Exclusively Distributed in the Philippines by:  
**STOLLER**  
Philippines, Inc.

Unit 801, Manila Luxury Condominium,  
Pearl Drive, Ortigas Center, Pasig City  
Philippines  
Phone: (632) 634-2121  
Web add: www.stoller.com.ph

## Harvest More 30 - 10 - 10

### DIRECTIONS FOR USE

The recommended time of application is early in the morning or late in the afternoon. Apply the lowest rate when leaves are young and higher rate to fully-grown leaves.

CROP	TIME OF APPLICATION	RATE (Tbsp/16L)
<b>RICE &amp; CORN</b>	Spray four times during the growing season 15 days after transplanting or 30 days after planting. If direct-seeded, repeat application every 2 weeks.	6-8
<b>Field &amp; Vegetable Crops:</b> Beans, Broccoli, Cabbage, Cauliflower, Celery, Cucumber, Lettuce, Onion, Melon, Potato, Pepper, Radish, Spinach, Squash, Tomato, etc.	Spray four times during the growing season 15 days after transplanting or 30 days after planting. If direct-seeded, repeat application every 2weeks	4-6
<b>Fruit &amp; Nut Crops:</b> Mango, Citrus, Grapes, Coffee, etc.	Apply early in the fruiting season or when improved vigor is desired. 3-4 applications may be done per season at 7-10 day intervals.  Caution: do not use nitrogenous fertilizer during the onset of the plant's reproductive stage.	4-6
<b>Ornamental Crops:</b> Orchids, Roses, Anthuriums, Chrysanthemums, Gladiolas, etc.	14 DAT up to flower formation	4-6

1 Tbsp = 10 grams

KEEP OUT OF REACH OF CHILDREN